



WINNICA TURNAU RONDO/REGENT 2021

WINE DESCRIPTION

The garnet colour reflects the richness of flavor and aromas carried by two red varieties of wine - rondo and regent. The nose presents the aroma of ripe cherries, dark fruits, smoke, leather and wood. Its concentration, bouquet and well-rounded tannins are obtained by long barrel ageing. Wine has potential for further development in the bottle.

WINE AROMAS

ripe cherry, smoked plum, spices

TASTE OF WINE

red berries, barrel, spices

FOODPAIRING

beef, cheese, game, grill

WINE PARAMETERS

residual sugar: 6,1 g/l

acidity: 5 g/l

alcohol content: 11,5%

GRAPE VARIETIES

2/3 rondo, 1/3 regent

VINIFICATION

Maceration 28 days on skins. Maturing 12 months in Slavonian oak barrique barrels. Barrels used once or twice.

AWARDS (ALL YEARS)

Notice: Array to string conversion in

`/var/www/winnicaturneau.pl/application/modules/default/views/scripts/wines/_catalog_en.phtml` on line 50

Array

Notice: Array to string conversion in

`/var/www/winnicaturneau.pl/application/modules/default/views/scripts/wines/_catalog_en.phtml` on line 50

Array

TERROIR

Winnica Turnau is located in Baniewice, in the southwestern part of the West Pomeranian Voivodeship. This unique region is influenced by air masses from the Atlantic Ocean and the Baltic Sea, as well as the proximity of the Oder River. All these factors shape a climate characterized by a long growing season, warm summers, and mild winters. Such conditions favor the cultivation of demanding grape varieties, allowing them to reach optimal ripeness while maintaining high acidity. Our region is distinguished by a rich network of oxbow lakes, gently rolling moraine hills, and an abundance of flora and fauna. The vineyard is situated on a moraine hill, where the predominant soils are sandy-clay, clay, moderately compact, and moderately permeable. This terroir, combined with ecological viticulture, results in grapes of exceptionally high quality.

WEATHER DURING THE GROWING PERIOD

Winter 2020/2021 was quite cold, with more snow cover than usual. The beginning of the year was relatively cool, and the rest of the year was characterized by great variability. Early spring arrived quickly, but March and April were cool with little rain. May was cold and rainy, with early spring frosts, but no losses. From hot and dry June the weather changed to a very rainy July, which slightly affected cluster formation. August surprised with changeable weather, and before harvest the rains threatened disease pressure. The autumn was warm with occasional showers and plenty of wind to dry the morning mists. The harvest began on time, and the first frosts began in November.